

aero® herbicides

AERO® line of Herbicides consists of a full range of non-selective weed killers, including oil and water-based products, liquids and granular, non-residuals and soil sterilants.

HIGHLIGHTS

- Non-selective, non-residual weed killer
- For terrestrial as well as aquatic uses
- Concentrated
- Fast-acting
- Deactivated on contact with soil


TYPICAL PROPERTIES

- pH: **6.0**
- Specific Gravity: **1.02**
- Diquat Dibromide: **1.85%**
- Appearance: **Black liquid**
- Odor: **Odorless**

SAFETY CAUTIONS

- Keep out of reach of children.
- May be fatal if swallowed, inhaled or absorbed through skin. Causes skin irritation. Do not get material on skin, eyes or clothing. Wear face shield, rubber gloves and rubber apron when handling concentrate.
- Refer to M.S.D.S. and product label for additional safety information and directions for use.

HMS RATINGS



Sudden Death

A Contact, Non-selective Vegetation Killer For The Control Of Broadleaf Weeds, Grasses And Aquatic Weeds

This herbicide when used as directed, will kill most forms of plant growth without damage to soil. Absorption and herbicidal action is usually quite rapid with effects visible in a few days.

GENERAL WEED CONTROL IN NON-CROP AREAS -

Controls broadleaf weeds, brush and grasses. Kills most annuals and top kills perennials. It kills through the leaves only. To kill above ground weed growth around buildings and other installations, walkways, patios, fence lines, parkways and driveways for control of undesirable broadleaved and grass-type weeds - 2½ gallons per 20 gallons of water. Apply for full coverage and thorough weed contact.

AQUATIC WEED CONTROL - Both surface and subsurface, in ponds, lakes and drainage ditches. Treat only 1/3 to ½ of the dense weed areas at a time and wait 14 days between treatments.

WATER TREATMENT

	ppm**
Algae (Spirogyra and Pithophora spp.)	½ to 1½
Bladderwort (Utricularia spp.)	¼ to ½
Coontail (Ceratophyllum Demersum)	1½ to 2½
Elodea (Elodea spp.)	1½ to 2½
Naiad (Naias spp.)	¼
Pondweed (Potamogeton spp.)	½
Watermilfoil (Myriophyllum spp.)	¼ to ½

**This product cation by weight. (See dilution chart.)

DILUTION CHART

Dept of Water and Gallons to Use per Acre at Indicated PPM Cation (by wt.)

Feet	1/4 ppm	½ ppm	1 ppm	1 ½ ppm
1	7 gal.	14 gal.	28 gal.	42 gal.
2	14 gal.	28 gal.	56 gal.	84 gal.
3	21 gal.	42 gal.	84 gal.	126 gal.
4	28 gal.	56 gal.	112 gal.	168 gal.
5	35 gal.	70 gal.	140 gal.	210 gal.

Foliage Spray Treatments: Amt./Acre

Salvinia (Salvinia rotundifolia)	10 to 15 gal.
Water hyacinth (Eichhornia crassipes)	10 to 15 gal.
Water lettuce (Pistia stratiotes)	10 to 15 gal.

FOLIAGE SPRAY TREATMENTS - Foliage spray treatments should be applied in sufficient water (150-200 gal. per acre) to thoroughly wet foliage.

See product label for complete directions.

distributed by:

aero®
CHEMICAL COMPANY

a division of:
ABC COMPOUNDING CO., INC.
Atlanta, Georgia Dallas, Texas